arXiv:math/0408402v1 [math.RA] 30 Aug 2004

Hochschild (co)homology dimension *

Yang Han

Institute of Systems Science, Academy of Mathematics and Systems Science,
Chinese Academy of Sciences, Beijing 100080, P.R.China.
E-mail: hany@mail.iss.ac.cn

Abstract

In 1989 Happel conjectured that for a finite-dimensional algebra A over an algebraically closed field k, gl.dim.$A < \infty$ if and only if hch.dim.$A < \infty$. Recently Buchweitz-Green-Madsen-Solberg gave a counterexample to Happel's conjecture. They found a family of pathological algebra A_q for which gl.dim.$A_q = \infty$ but hch.dim.$A_q = 2$. These algebras are pathological in many aspects, however their Hochschild homology behaviors are not pathological any more, indeed one has hh.dim.$A_q = \infty =$ gl.dim.A_q. This suggests to pose a seemingly more reasonable conjecture by replacing Hochschild cohomology dimension in Happel's conjecture with Hochschild homology dimension: gl.dim.$A < \infty$ if and only if hh.dim.$A < \infty$ if and only if hh.dim.$A = 0$. The conjecture holds for commutative algebras and monomial algebras. In case A is a truncated quiver algebras these conditions are equivalent to the quiver of A has no oriented cycles. Moreover, an algorithm for computing the Hochschild homology of any monomial algebra is provided. Thus the cyclic homology of any monomial algebra can be read off in case the underlying field is characteristic 0.

Mathematics Subject Classification (2000): 16E40, 16E10, 16G10

Let k be a fixed field. All algebras considered here are of the form $A = kQ/I$ where Q is a finite quiver and I is an admissible ideal of the path algebra kQ. We refer to [ARS] for the theory of quivers and their representations. It is well-known that in case k is an algebraically closed field, up to Morita

*Project 10201004 supported by NSFC.

equivalence, any finite-dimensional k-algebra can be written as this form (cf. [G1]).

1. Hochschild (co)homology

In this section we recall some definitions and properties on Hochschild (co)homology.

(1.1) The complex $C_*(A) = (A^{\otimes *+1}, b_*) : \cdots \to A^{\otimes n+1} \xrightarrow{b_n} A^{\otimes n} \to \cdots \to A^{\otimes 2} \xrightarrow{b_1} A \to 0$ with $b_n(a_0, a_1, ..., a_n) := \sum_{i=0}^{n-1}(-1)^i(a_0, ..., a_i a_{i+1}, ..., a_n) + (-1)^n(a_n a_0, a_1, ..., a_{n-1})$, where $(a_0, ..., a_n) := a_0 \otimes \cdots \otimes a_n$ and $\otimes := \otimes_k$, is called the *Hochschild complex* or *cyclic bar complex* of A.

The homology group $HH_n(A) := H_n(C_*(A))$ is called the nth *Hochschild homology group* of A (cf. [Lod]). Clearly $HH_n(A)$ is a k-vector space, we denote $\dim_k HH_n(A)$ by $hh_n(A)$.

(1.2) The cohomology group of the complex $C^*(A) = (\mathrm{Hom}_k(A^{\otimes *}, A), d^*) : 0 \to A \xrightarrow{d^0} \mathrm{Hom}_k(A, A) \xrightarrow{d^1} \cdots \to \mathrm{Hom}_k(A^{\otimes n}, A) \xrightarrow{d^n} \mathrm{Hom}_k(A^{\otimes n+1}, A) \to \cdots$ with $(d^n(f))(a_0, a_1, ..., a_n) := a_0 f(a_1, ..., a_n) + \sum_{i=0}^{n-1}(-1)^{i+1} f(a_0, ..., a_i a_{i+1}, ..., a_n) + (-1)^{n+1} f(a_0, ..., a_{n-1})a_n$, $HH^n(A) := H^n(C^*(A))$, is called the nth *Hochschild cohomology group* of A (cf. [Ho]). Clearly $HH^n(A)$ is a k-vector space, we denote $\dim_k HH^n(A)$ by $hh^n(A)$.

(1.3) The chain complex $C'_*(A) = (A^{\otimes *+2}, b'_*) : \cdots \to A^{\otimes n+2} \xrightarrow{b'_n} A^{\otimes n+1} \to \cdots \to A^{\otimes 3} \xrightarrow{b'_1} A^{\otimes 2} \to 0$ with $b'_n(a_0, a_1, ..., a_{n+1}) := \sum_{i=0}^{n}(-1)^i(a_0, ..., a_i a_{i+1}, ..., a_n)$ is called the *bar complex* or *standard complex* of A.

The bar complex $C'_*(A)$ is a projective resolution of A as an A^e-module where $A^e := A \otimes A^{op}$. Since $A \otimes_{A^e} C'_*(A) = C_*(A)$ and $\mathrm{Hom}_{A^e}(C'_*(A), A) = C^*(A)$, we have $HH_n(A) = \mathrm{Tor}_n^{A^e}(A, A)$ and $HH^n(A) = \mathrm{Ext}^n_{A^e}(A, A)$. These two formulae are very useful to calculate Hochschild (co)homology by constructing a convenient projective resolution of A over A^e.

(1.4) Lemma. *Let A and B be two algebras.*

(1) $HH_n(\left[\begin{smallmatrix} A & {}_AM_B \\ 0 & B \end{smallmatrix}\right]) = HH_n(A) \oplus HH_n(B)$ *for any A-B-bimodule M. In particular $HH_n(A \times B) = HH_n(A) \oplus HH_n(B)$.*

(2) *If A and B are derived equivalent then $HH_n(A) = HH_n(B)$. In particular iterate tilting and Morita equivalence preserves Hochschild homology.*

(3) $HH_n(A \otimes B) = \oplus_{i+j=n} HH_i(A) \otimes HH_j(B)$.

Proof. (1) [Lod; Theorem 1.2.15]. (2) [K1; Theorem 2.2]. (3) [Mac; Chap.VIII, Theorem 7.4]. □

(1.5) Lemma. *Let A and B be two algebras.*

(1) $HH^n(A \times B) = HH^n(A) \oplus HH^n(B)$.

(2) *If A and B are derived equivalent then $HH^n(A) = HH^n(B)$. In particular iterate tilting and Morita equivalence preserves Hochschild cohomology.*

(3) $HH^n(A \otimes B) = \oplus_{i+j=n} HH^i(A) \otimes HH^j(B)$.

Proof. (1) [CE; Chap.IX, Theorem 5.3]. (2) [R; Proposition 2.5]. (3) [Mac; Chap.VIII, Theorem 7.4]. □

Remark. For Hochschild cohomology in general we have no similar formula to Lemma (1.4) (1). An obvious counterexample is the Kronecker algebra $\left[\begin{smallmatrix} k & k^2 \\ 0 & k \end{smallmatrix}\right]$. Its Hochschild cohomology groups can be obtained by [Ha; Proposition 1.6]. The relation between the triangular matrix algebra $\left[\begin{smallmatrix} A & {}_AM_B \\ 0 & B \end{smallmatrix}\right]$ and A, B was observed in [MiP].

2. Hochschild (co)homology dimension

In this section we give the definitions and some properties of Hochschild (co)homology dimension.

(2.1) Definition. ([B; Definition 3.1]) The *Hochschild homology dimension* of A is $\mathrm{hh.dim.}A := \inf\{n \in \mathbb{N}_0 | hh_i(A) = 0 \text{ for } i > n\}$.

Proposition. *Let A and B be two algebras.*

(1) $\mathrm{hh.dim.}\left[\begin{smallmatrix} A & {}_AM_B \\ 0 & B \end{smallmatrix}\right] = \max\{\mathrm{hh.dim.}A, \mathrm{hh.dim.}B\}$ *for any A-B-bimodule M. In particular* $\mathrm{hh.dim.}(A \times B) = \max\{\mathrm{hh.dim.}A, \mathrm{hh.dim.}B\}$.

(2) *If A and B are derived equivalent then* $\mathrm{hh.dim.}A = \mathrm{hh.dim.}B$. *In particular iterate tilting and Morita equivalence preserves Hochschild homology dimension.*

(3) $\mathrm{hh.dim.}(A \otimes B) = \mathrm{hh.dim.}A + \mathrm{hh.dim.}B$.

Proof. It follows from Lemma (1.4) □

(2.2) Definition. The *Hochschild cohomology dimension* of A is $\mathrm{hch.dim.}A := \inf\{n \in \mathbb{N}_0 | hh^i(A) = 0 \text{ for } i > n\}$.

Proposition. *Let A and B be two algebras.*

(1) $\mathrm{hch.dim.}(A \times B) = \max\{\mathrm{hch.dim.}A, \mathrm{hch.dim.}B\}$.

(2) *If A and B are derived equivalent then* $\mathrm{hch.dim.}A = \mathrm{hch.dim.}B$. *In particular iterate tilting and Morita equivalence preserves Hochschild cohomology dimension.*

(3) $\mathrm{hch.dim.}(A \otimes B) = \mathrm{hch.dim.}A + \mathrm{hch.dim.}B$.

Proof. It follows from Lemma (1.5). □

Remark. For some special class of algebras we have some other nice properties on Hochschild cohomology dimension: If A is hereditary then hch.dim.$A \leq 1$, in this case hch.dim.$A = 0$ if and only if Q is a tree (cf. [Ha; Proposition 1.6 and Corollary 1.6]). If A is radical square zero then hch.dim.$A = 0$ if and only if Q is a tree (cf. [Ha; Proposition 2.3]).

3. Relation with global dimension

In this section we consider the relation between the Hochschild (co) homology dimension and the global dimension of an algebra. This is the main part of the paper.

(3.1) Conjecture. ([Ha]) hch.dim.$A < \infty$ *if and only if* gl.dim.$A < \infty$.

Let $Q = (Q_0, Q_1)$ where Q_0 (resp. Q_1) is the set of vertices (resp. arrows) in Q. For any arrow $a \in Q_1$ denote by $s(a)$ and $e(a)$ the starting point and the ending point of a respectively. A path $a_1 \cdots a_n$ with $n \geq 1$ is called an *oriented cycle* or a *cycle* of length n if $e(a_i) = s(a_{i+1})$ for $i = 1, ..., n-1$ and $e(a_n) = s(a_1)$. Happel's conjecture holds for *truncated quiver algebras*, i.e., the algebras kQ/k^nQ where $n \geq 2$ and k^nQ denotes the ideal of kQ generated by all paths of length $\geq n$ (cf. [Ci2]).

Proposition. ([Loc; Theorem 3]) *Let A be a truncated quiver algebra. Then the following are equivalent:*

(1) hch.dim.$A < \infty$;

(2) gl.dim.$A < \infty$;

(3) *The quiver of A has no oriented cycles.*

However Happel's conjecture is not true in general.

(3.2) Pathological algebras. Let $A := A_q := k\langle x, y\rangle/(x^2, xy + qyx, y^2)$ with $q \in k$. The algebra A is very pathological for some special q. It is used to construct counterexamples for a few problems:

(1) ([BGMS]) By constructing a minimal projective resolution of A over A^e Buchweitz-Green-Madsen-Solberg show that hch.dim.$A = 2$ in the case q is neither zero nor a root of unity, but gl.dim.$A = \infty$. This provides a negative answer to Happel's conjecture.

(2) ([Sc]) By constructing a non-projective A_q-module without self-extension for some special q, Schulz gives a negative answer to Tachikawa's conjecture: Over a quasi-Frobenius algebra, a finitely generated module without self-extensions is projective.

(3) ([LS]) In case q is not a root of unity, the trivial extension of A_q is a local symmetric algebra whose AR-quiver contains a bounded infinite DTr-orbit. Thus Liu-Schulz give a negative answer to Ringel's problem: whether

the number of modules having the same length in a connected component of the AR-quiver is always finite.

(4) ([SY]) Richard showed that A_q and $A_{q'}$ are socle equivalent self-injective algebras which are not stably equivalent in case $q' \neq q$ or q^{-1}.

Though these algebras are quite pathological in many aspects, their Hochschild homology behaviors are not pathological any more.

Proposition. *For any* $q \in k$, hh.dim.$A = \infty =$ gl.dim.A.

Proof. Recall the minimal projective resolution constructed in [BGMS]:

For any $n \geq 0$ construct $\{f_i^n\}_{i=0}^n \subseteq A^{\otimes n}$: First $\{f_0^0\} \subseteq A^{\otimes 0} = k$ with $f_0^0 := 1$. Second $f_0^1 := x$ and $f_1^1 := y$. Third $f_{-1}^n := 0 =: f_{n+1}^n$, and for any $n \geq 2$, $f_i^n := f_{i-1}^{n-1} \otimes y + q^i f_i^{n-1} \otimes x$ with $i \in \{0, 1, ..., n\}$. Here A is identified with $k \otimes_k A$. Note that $f_i^n = x \otimes f_i^{n-1} + q^{n-i} y \otimes f_{i-1}^{n-1}$.

Let $P^n := \oplus_{i=0}^n A \otimes_k f_i^n \otimes_k A \subseteq A^{\otimes(n+2)}$ for all $n \geq 0$. Set $\tilde{f}_i^n := 1 \otimes f_i^n \otimes 1$ for $n \geq 1$ and $\tilde{f}_0^0 := 1 \otimes 1$. Define $\delta^n : P^n \to P^{n-1}$ by letting $\delta^n(\tilde{f}_i^n) := x\tilde{f}_i^{n-1} + q^{n-i} y \tilde{f}_{i-1}^{n-1} + (-1)^n \tilde{f}_{i-1}^{n-1} y + (-1)^n q^i \tilde{f}_i^{n-1} x$. Then (P^*, δ^*) is a minimal projective resolution of A over A^e.

Applying the functor $A \otimes_{A^e} -$ to the minimal projective resolution we have $A \otimes_{A^e} (P^*, \delta^*) = (\oplus_{i=0}^* A \otimes_k f_i^*, \tau^*)$ where τ^n is defined by $\tau^n(\lambda \otimes f_i^n) = \lambda x \otimes f_i^{n-1} + (-1)^n q^i x\lambda \otimes f_i^{n-1} + q^{n-i} \lambda y \otimes f_{i-1}^{n-1} + (-1)^n y\lambda \otimes f_{i-1}^{n-1}$.

Note that A has a basis $\{1, y, x, yx\}$. Thus $A \otimes_k f_i^n$ has a basis $\{1 \otimes f_i^n, y \otimes f_i^n, x \otimes f_i^n, yx \otimes f_i^n\}$. Furthermore $\oplus_{i=0}^n A \otimes_k f_i^n$ has a basis $\{1 \otimes f_i^n, y \otimes f_i^n, x \otimes f_i^n, yx \otimes f_i^n\}_{i=0}^n$. If $n \geq 1$ then $\oplus_{i=0}^{n-1} A \otimes_k f_i^{n-1}$ has a basis $\{1 \otimes f_i^{n-1}, y \otimes f_i^{n-1}, x \otimes f_i^{n-1}, yx \otimes f_i^{n-1}\}_{i=0}^{n-1}$. Moreover $\tau^n(1 \otimes f_i^n) = (1 + (-1)^n q^i)(x \otimes f_i^{n-1}) + (q^{n-i} + (-1)^n)(y \otimes f_{i-1}^{n-1})$, $\tau^n(y \otimes f_i^n) = (1 + (-1)^{n+1} q^{i+1})(yx \otimes f_i^{n-1})$, $\tau^n(x \otimes f_i^n) = (-q^{n-i+1} + (-1)^n)(yx \otimes f_{i-1}^{n-1})$ and $\tau^n(yx \otimes f_i^n) = 0$. Write τ^n as a matrix, it is not difficult to see that $\mathrm{rank}(\tau^n) \leq \begin{cases} 2n-1, & \text{if } n \text{ odd;} \\ 2n+1, & \text{if } n \text{ even.} \end{cases}$ Thus $\mathrm{rank}(\tau^{n+1}) \leq \begin{cases} 2n+3, & \text{if } n \text{ odd;} \\ 2n+1, & n \text{ even.} \end{cases}$ Therefore $hh_n(A) = \dim_k \mathrm{Ker}(\tau^n) - \dim_k \mathrm{Im}(\tau^{n+1}) = \dim_k P^n - \dim_k \mathrm{Im}(\tau^n) - \dim_k \mathrm{Im}(\tau^{n+1}) = 4(n+1) - \mathrm{rank}(\tau^n) - \mathrm{rank}(\tau^{n+1}) \geq 4(n+1) - (4n+2) = 2$. Indeed $hh_n(A) \geq 2$ for all n since $hh_0(A) = \dim_k A/[A, A] = \begin{cases} 4, & \text{if } q = 1; \\ 3, & \text{if } q \neq 1. \end{cases}$ □

(3.3) Proposition (3.2) suggests us to pose a seemingly more reasonable conjecture by replacing hch.dim.A in Happel's conjecture with hh.dim.A. Indeed the following result, which is essentially due to Keller, makes us expect more. Denote by E the maximal semisimple subalgebra of A, i.e., $E := kQ_0$, and by r the Jacobson radical of A. Then one has $A = E \oplus r$.

Proposition. *If gl.dim.$A < \infty$ then* $hh_i(A) = \begin{cases} |Q_0|, & \text{if } i = 0; \\ 0, & \text{if } i \geq 0. \end{cases}$

Proof. If gl.dim.$A < \infty$ then by [K2; Proposition 2.5] we have

$HC_n(A, \mathrm{rad}A) = 0$ for all $n \geq 0$. Using the long exact sequence

$$\cdots \to HH_n(A,r) \to HC_n(A,r) \to HC_{n-2}(A,r) \to \cdots$$

$$\to HH_2(A,r) \to HC_2(A,r) \to HC_0(A,r) \to HH_1(A,r)$$
$$\to HC_1(A,r) \to 0 \to HH_0(A,r) \to HC_0(A,r) \to 0$$

(cf. [Lod; E2.2.4]) we have $HH_n(A,r) = 0$ for all $n \geq 0$. Using the long exact sequence

$$\cdots \to HH_n(A,r) \to HH_n(A) \to HH_n(E) \to$$

$$\cdots \to HH_0(A,r) \to HH_0(A) \to HH_0(E) \to 0$$

(cf. [Lod; §1.1.16]), we have $HH_n(A) = HH_n(E)$ for all $n \geq 0$. Thus the proposition follows from $hh_i(E) = \left\{ \begin{smallmatrix} |Q_0| & \text{if } i = 0; \\ 0 & \text{if } i \geq 0. \end{smallmatrix} \right.$ (cf. [Lod; E1.1.1]). □

Corollary. *If* gl.dim.$A < \infty$ *then* hh.dim.$A = 0$.

(3.4) Conjecture. *The following conditions are equivalent:*
(1) hh.dim.$A < \infty$;
(2) hh.dim.$A = 0$;
(3) gl.dim.$A < \infty$.

(3.5) Now we show that Conjecture (3.4) holds for commutative algebras. We refer to [Mat] for commutative algebra theory. The following result is essentially due to Avramov and Vigueé-Poirrier.

Theorem. *Let A be a commutative algebra. Then the following conditions are equivalent:*
(1) hh.dim.$A < \infty$;
(2) hh.dim.$A = 0$;
(3) gl.dim.$A < \infty$
(4) gl.dim.$A = 0$.

Proof. By Corollary (3.1) it is enough to show that (1) $\Rightarrow$ (4). Since A is commutative, it is the direct product of local algebras. Thus without loss of generality we may assume that A is local. If hh.dim.$A < \infty$ then by [AV; Theorem] A is smooth. By [Mat; p.216, Lemma 1], A is regular. Furthermore gl.dim.$A = 0$ by Serre's theorem (cf. [Mat; Theorem 19.2]). □

(3.6) Now we prove that Conjecture (3.4) holds for *monomial algebras* or *zero relation algebras* in some literature, i.e., the algebras $A = kQ/I$ with I generated by some paths of length ≥ 2.

A cycle $a_1 \cdots a_n$ with $\{a_1, ..., a_n\} \subseteq Q_0$ is said to be *basic* if the vertices $s(a_1), ..., s(a_n)$ are distinct from each other (cf. [LZ]). A cycle is said to be

proper if it is not a power (≥ 2) of any cycle (cf. [Ci3]). Note that proper cycles are called basic cycles in [Sk]. Denote by Q_n^c (resp. Q_n^p) the set of cycles (resp. proper cycles) of length n. The truncated quiver algebra whose quiver is a basic cycle is called a *truncated basic cycle algebra*. The Hochschild homology of a truncated basic cycle is known (cf. [Sk] and [Z]). If A is a monomial algebra then A can be naturally defined to be an $\mathbb{N}_0$-graded algebra by using the length of the paths. Thus the i-th Hochschild homology group $HH_n(A)$ becomes an $\mathbb{N}_0$-graded k-vector space by $\deg_{\mathbb{N}_0} a_0 \otimes \cdots \otimes a_n := \sum_{i=0}^n \deg_{\mathbb{N}_0} a_i$ for $a_i \in A, i = 0, 1, ..., n$. The degree j part of $HH_i(A)$ is denoted by $HH_{i,j}(A)$. Let C_q be the cyclic group of order q. Then C_q acts on Q_q^c by $g \cdot (a_1 \cdots a_{q-1} a_q) := (a_q a_1 \cdots a_{q-1})$. Under this action the orbit of the cycle Y is denoted by O_Y.

Proposition. ([Sk; Theorem 2]) *Let $A = kQ/k^nQ$ be a truncated quiver algebra and $q = cn + e$ for $0 \leq e \leq n-1$. Then $HH_{p,q}(A) =$*

$$\begin{cases} k^{a_q}, & \text{if } 1 \leq e \leq n-1 \text{ and } 2c \leq p \leq 2c+1; \\ \oplus_{r|q}(k^{(n,r)-1} \oplus \mathrm{Ker}(\cdot \frac{n}{(n,r)} : k \to k))^{b_r}, & \text{if } e = 0 \text{ and } 0 < 2c-1 = p; \\ \oplus_{r|q}(k^{(n,r)-1} \oplus \mathrm{Coker}(\cdot \frac{n}{(n,r)} : k \to k))^{b_r}, & \text{if } e = 0 \text{ and } 0 < 2c = p; \\ k^{|Q_0|}, & \text{if } p = q = 0; \\ 0, & \text{otherwise.} \end{cases}$$

Here $a_q := \mathrm{card}(Q_q^c/C_q), (n,r) := \gcd(n,r)$ and $b_r := \mathrm{card}(Q_r^p/C_r)$.

Corollary 1. *Let $A = kQ/k^nQ$ where Q is a basic cycle of length l. Then*

$$hh_p(A) = \begin{cases} l + [\frac{n-1}{l}], & \text{if } p = 0; \\ \left[\frac{[\frac{p}{2}]n+n-1}{l}\right] - \left[\frac{[\frac{p}{2}]n}{l}\right], & \text{if } (p \geq 1) \wedge (l \nmid [\frac{p+1}{2}]n); \\ \left[\frac{[\frac{p}{2}]n+n-1}{l}\right] - \left[\frac{[\frac{p}{2}]n}{l}\right] + (n,l) - 1, & \text{if } (p \geq 1) \wedge (l | [\frac{p+1}{2}]n) \wedge (\mathrm{char} k \nmid \frac{n}{(n,l)}); \\ \left[\frac{[\frac{p}{2}]n+n-1}{l}\right] - \left[\frac{[\frac{p}{2}]n}{l}\right] + (n,l), & \text{if } (p \geq 1) \wedge (l | [\frac{p+1}{2}]n) \wedge (\mathrm{char} k | \frac{n}{(n,l)}). \end{cases}$$

Here $[x]$ denotes the largest integer i satisfying $i \leq x$.

Corollary 2 *If $A = kQ/k^nQ$ where Q is a basic cycle of length l. Then* $\mathrm{hh.dim}.A = \infty$.

Proof. If $(n,l) \geq 2$ then for any $p = 2ml - 1$ with $m \geq 1$ one has $hh_p(A) \geq 1$. If $(n,l) = 1$ and $l \leq n-1$ then $\left[\frac{[\frac{p}{2}]n+n-1}{l}\right] > \left[\frac{[\frac{p}{2}]n}{l}\right]$ for any $p \geq 1$. Thus $hh_p(A) \geq 1$ for any $p \geq 1$. If $(n,l) = 1$ and $l > n-1$ then there are integers u and v such that $un + vl = 1$. Thus for any $p = 2(ml + u - 1)$ with $m \gg 0$ one has $\left[\frac{[\frac{p}{2}]n+n-1}{l}\right] = \left[\frac{[\frac{p}{2}]n+n-un-vl}{l}\right] = mn - v$ and $\left[\frac{[\frac{p}{2}]n}{l}\right] < mn - v$. Therefore $hh_p(A) \geq 1$. $\square$

Let $A = kQ/I$ be a monomial algebra. Let $C' = a_1 \cdots a_n$ be a cycle in Q. Then there is a basic cycle quiver C with $|C_0| = n$ and a natural quiver

morphism $C \to C' \to Q$. We define the relations on C by pulling back the relations in Q, i.e., a path in C is a relation if and only if its image in Q is a relation. The algebra Z given by quiver C and the relations is called a *cycle algebra overlying* A. A cycle algebra Z overlying A is said to be *minimal* if Z is not a covering of a smaller cycle algebra overlying A (cf. [IZ]). For covering theory we refer to [G2] and [MarP]. Clearly minimal cycle algebras overlying A are just those cycle algebras overlying A constructed from proper cycles in Q. Denote by $HC_n(A)$ the n-th cyclic homology group of A (cf. [Lod]).

Theorem 1. *Let A be a monomial algebra. Then $HH_n(A) \cong \oplus_Z HH_n(Z)$ and $HC_n(A) \cong \oplus_Z HC_n(Z)$ where the index Z runs through all minimal cycle algebra over A.*

Proof. By [Ci3; Lemma 2.3] the Hochschild homology and the cyclic homology of A is just the Hochschild homology and the cyclic homology of the *E-normalized mixed complex* $\bar{C}_E(A) = (A \otimes_{E^e} (r^{\otimes_E n}), b, B)$ with b the Hochschild boundary and $B : A \otimes_{E^e} (r^{\otimes_E n}) \to A \otimes_{E^e} (r^{\otimes_E n+1})$ given by the formula $B(a_0, a_1, ..., a_n) = \sum_{i=0}^{n} (-1)^{in}(1, a_i, ..., a_n, (a_0)_r, ..., a_{i-1})$, where $a = a_E + a_r$ is the decomposition of an element a of A in the direct sum $A = E \oplus r$. Note that the k-vector space $A \otimes_{E^e} (r^{\otimes_E n}) = \oplus_{a_0, r_1, ..., r_n} k(a_0 \otimes r_1 \otimes \cdots \otimes r_n)$ (where $a_0, r_1, ..., r_n$ runs all paths sequences with $a_0 r_1 \cdots r_n$ being a cycle) $= \oplus_{\text{orbit } O_Y \text{ of proper cycle } Y \text{ in } Q} \oplus_{a_0, r_1, ..., r_n} k(a_0 \otimes r_1 \otimes \cdots \otimes r_n)$ (where $a_0, r_1, ..., r_n$ runs all paths sequences with $a_0 r_1 \cdots r_n = X^m$ for some $X \in O_Y$ and m) $= \oplus_{\text{minimal cycle algebra } Z \text{ overlying } A} Z \otimes_{F^e} (s^{\otimes_F n})$ where F is the maximal semisimple subalgebra of Z and s is the Jacobson radical of Z. Clearly this induces an isomorphism of mixed complex from $\bar{C}_E(A)$ to $\oplus_Z \bar{C}_F(Z)$ where Z runs through all minimal cycle algebras over A. By [Lod; Proposition 2.5.15] we are done. □

Algorithm. The Hochschild homology of any monomial algebra A can be computed as following:

(1) Determine the set $\mathcal{O}$ of the orbits of proper cycles in the quiver of A.

(2) Determine the set $\mathcal{M}$ of the minimal cycle algebras overlying A corresponding to the elements in $\mathcal{O}$.

(3) Apply the reduction of dropping projective modules in [IZ; p. 512], [IZ; Theorem 2.3] and (3.6) Corollary 1 to obtain $HH_n(Z)$ for each $Z \in \mathcal{M}$ and for all $n \geq 0$.

(4) Apply (3.6) Theorem 1 to obtain $HH_n(A)$ for all $n \geq 0$.

In case char$k = 0$, the cyclic homology of A can be read off:

(5) Apply [Lod; Theorem 4.1.13] to obtain $HC_n(A)$.

Theorem 2. *Let A be a monomial algebra. Then the following conditions are equivalent:*

(1) hh.dim.$A < \infty$;
(2) hh.dim.$A = 0$;
(3) gl.dim.$A < \infty$.

Proof. By (3.1) Corollary 1 it is enough to show that (1) $\Rightarrow$ (3): If gl.dim.$A = \infty$ then by [IZ; Proposition 3.2] there is a minimal cycle algebra Z with gl.dim.$Z = \infty$. Via a series of dropping projective modules, Z can be reduced to a nonsemisimple truncated basic cycle algebra Z' (cf. [IZ; p. 512]). By (3.6) Corollary 2 and [IZ; Theorem 2.3], one has hh.dim.Z = hh.dim.$Z' = \infty$. Thus by (3.6) Theorem 1 one has hh.dim.$A \geq$ hh.dim.$Z = \infty$. It is a contradiction. □

Remark 1. The equivalence between (2) and (3) is the main result of [IZ].

It is well-known that the algebras whose Gabriel quiver have no oriented cycles are of finite global dimension (cf. [ENN; Corollary 6]) and Hochschild homology dimension 0 (cf. [Ci1]). We show in the following that their converses hold for truncated quiver algebras.

Theorem 3. *Let A be a truncated quiver algebra (in particular a radical square zero algebra). Then the following conditions are equivalent:*
(1) hh.dim.$A < \infty$;
(2) hh.dim.$A = 0$;
(3) gl.dim.$A < \infty$;
(4) *Q has no oriented cycles.*

Proof. (4) $\Rightarrow$ (3): It is well-known (cf. [ENN; Corollary 6]). By (3.1) Corollary 1 it is enough to show that (1) $\Rightarrow$ (4): If Q has an oriented cycle then we take the shortest cycle Q' in Q which must be a basic cycle. Let $I' = I \cap kQ'$. Then $Z := kQ'/I'$ is an truncated basic cycle algebra. By [LZ; Theorem 2] or (3.6) Theorem 1, $HH_n(Z)$ is a direct summand of $HH_n(A)$ for all $n \geq 0$. By (3.6) Corollary 2 we have hh.dim.$Z = \infty$ which implies hh.dim.$A = \infty$. It is a contradiction. □

Remark 2. In general the finiteness of the global dimension does not imply that the Gabriel quiver of the algebra has no oriented cycle. The simplest counterexample is the algebra $A = kQ/I$ where $Q_0 = \{1, 2\}$, $Q_1 = \{\alpha : 1 \to 2, \beta : 2 \to 1\}$ and $I = \langle \beta\alpha \rangle$. Clearly gl.dim.$A = 2$.

References

[ARS] M. Auslander, I. Reiten and S.O. SmalØ, Representation theory of artin algebras, Cambridge studies in advanced mathematics 36, Cambridge university press, Cambridge, 1995.

[AV] L.L. Avramov and M. Vigueé-Poirrier, Hochschild homology criteria for smoothness, Internat. Math. Research Notices 1 (1992), 17-25.

[B] J.L. Block, Cyclic homology of filtered algebras, K-theory 1 (1987), 515–518.

[BGMS] R.O. Buchweitz, E.L. Green, D. Madsen and Ø. Solberg, Finite Hochschild cohomology without finite global dimension, Preprint 2004.

[CE] H. Cartan and S. Eilenberg, Homological algebra, Princeton University Press, 1956.

[Ci1] C. Cibils, Hochschild homology of an algebra whose quiver has no oriented cycles, Springer Lecture Notes in Math. 1177, 1986, 55-59.

[Ci2] C. Cibils, Rigidity of truncated quiver algebras, Adv. Math. 79 (1990), 18-42.

[Ci3] C. Cibils, Cyclic and Hochschild homology of 2-nilpotent algebras, K-theory 4 (1990), 131–141.

[ENN] S. Eilenberg, H. Nagao and T. Nakayama, On the dimension of modules and algebras, IV, Dimension of residue rings of hereditary rings, Nagoya Math. J. 8 (1955), 49–57.

[G1] P. Gabriel, Auslander-Reiten sequences and representation- finite algebras, In: Representation Theory I, Springer Lecture Notes in Math. 831, 1980, 1–71.

[G2] P. Gabriel, The universal cover of a representation-finite algebra, Springer Lecture Notes in Math. 903, 1980, 68–105.

[Ha] D. Happel, Hochschild cohomology of finite-dimensional algebras, Springer Lecture Notes in Math. 1404, 1989, 108–126.

[Ho] G. Hochschild, On the cohomology groups of an associate algebra, Ann. Math. 46 (1945), 58-67.

[IZ] K. Igusa and D. Zacharia, On the cyclic homology of monomial relation algebras, J. Algebra 151 (1992), 502–521.

[K1] B. Keller, Invariance of cyclic homology under derived equivalence, Can. Math. Soc. Conf. Proc. Vol. 18, 1996, 353-361.

[K2] B. Keller, Invariance and localization for cyclic homology of DG algebras, J. Pure Appl. Algebra 123 (1998), 223–273.

[LS] S.P. Liu and R. Schulz, The existence of bounded infinite DTr-orbits, Proc. Amer. Math. Soc. 122 (1994), 1003-1005.

[LZ] S.X. Liu and P. Zhang, Hochschild homology of truncated algebras, Bull. London Math. Soc. 26(1994), 427–430.

[Loc] A.C. Locateli, Hochschild cohomology of truncated quiver algebras, Comm. Algebra 27 (1999), 645-664.

[Lod] J.L. Loday, Cyclic homology, Grundlehren 301, Springer, Berlin, 1992.

[Mac] S. Mac Lane, Homology, Grundlehren 114, Third corrected printing, Springer-Verlag, 1975.

[MarP] R. Martinez-Villa and J.A. de la Peña, The universal cover of a quiver with relations, J. Pure Appl. Algebra 30 (1983), 277–292.

[Mat] H. Matsumura, Commutative ring theory, Translated by M. Reid, Cambridge Univ. Press, 1986.

[MiP] S. Michelena and M. Platzeck, Hochschild cohomology of triangular matrix algebras, J. Algebra 233 (2000), 502-525.

[R] J. Richard, Derived equivalences as derived functors, J. London Math. Soc. 43 (1991), 37-48.

[Sc] R. Schulz, A non-projective module without self-extensions, Arch. Math. (Basel) 62 (1994), 497-500.

[Sk] E. Sköldberg, The Hochschild homology of truncated and quadratic monomial algebras, J. London Math. Soc. 59 (1999), 76–86.

[SY] A. Skowroński and K. Yamagata, Socle deformations of self-injective algebras, J. London Math. Soc. 72 (1996), 545-566.

[Z] P. Zhang, The Hochschild homology of a truncated basic cycle, Algebra Colloq. 5 (1998), 77-84.